
Metallic bonding

Metals LOSE ELECTRONS to form POSITIVE IONS




GIANT structures of atoms in a REGULAR pattern

Delocalised electrons are free to move.

What is a metallic bond?

Sharing delocalised electrons - STRONG metallic bonds.

Which type of bonding is it?



Ionic bonding

Metals LOSE ELECTRONS to form **POSITIVE IONS** Non-metals GAIN ELECTRONS to form NEGATIVE IONS

Electrons transferred from metal to non-metal

Ions have electronic structure of a noble gas

What is an ionic bond? STRONG electrostatic force of attraction between oppositely charged ions

How do we quickly work out the charges on ions?

Group	Electrons in outer shell	Charge on ion
1	1	1+
2	2	2+
6	6	2-
7	7	1-

L13 – 19: Structure and Bonding

Covalent **Bonding**

Two non-metals will SHARE pairs of electrons


STRONG bond formed.





small molecules

A small group of atoms sharing electrons











Giant Structures

Many atoms sharing electrons


















Limitations of these models

Model	Limitations
Dot and cross	Looks like electrons aren't identical Electrons look like they are in fixed positions
н-N-H Displayed formula	Doesn't show true shape of the molecule
Ball and stick	Can attempt to show 3D shape but doesn't show electrons

Properties of Metallic **Substances**

Metals have high melting and boiling points because...

...they are giant structures of atoms with strong metallic bonding

Can be bent or shaped because...

...atoms are arranged in LAYERS which can **SLIDE** over each



Allovs are harder than pure metals because...

Allovs are a mixture of two or more elements, at least one of which is a metal

...the layers are **DISTORTED** so can't slide over each other

Metals are good conductors of electricity and thermal energy Because...

...the electrons are free to move and carry thermal energy and charge

Properties of Ionic Substances

Ionic compounds have high melting and boiling points because...



...they are giant structures of atoms (giant ionic lattice) with strong electrostatic forces of attraction in ALL **DIRECTIONS** between oppositely charged ions.

A large amount of energy is needed to break the many strong bonds.

Only conduct electricity when melted or dissolved in water because...


...the ions are free to move and so charge can flow.

As ionic compounds are made of CHARGED IONS, they can CONDUCT ELECTRICITY but ONLY if the ions can MOVE.

Structure and Bonding

Small molecules

Small molecules have relatively low melting and boiling points because...



...intermolecular forces are overcome on melting and boiling and these are weak forces.

The bigger the size of the molecule the higher the melting and boiling point because...

...intermolecular forces increase with the size of the molecules.

Don't conduct electricity because...

...the molecules have no overall electric charge.

Properties of Covalent substances



Polymers are solids at room temperature

because...

...intermolecular forces increase with the size of the molecules and polymer molecules are very large.

Diamond is very hard, has a very high melting and boiling point and doesn't conduct electricity

because...

...each carbon is bonded to 4 other carbons by strong covalent bonds. There are no free electrons.

Graphite is very hard, has a very high melting and boiling point and does conduct electricity


because.. in the layer

Giant covalent compounds have high melting and boiling points

because...

...all of the atoms linked by strong covalent bonds.

Graphene is strong, light and an excellent conductor of thermal energy and electricity because...



.it is a single layer of graphite so has free electrons.

> Fullerenes (e.g. carbon nanotubes) are extremely strong and are excellent conductors of thermal energy and electricity because...

...each carbon is bonded to 3 other carbons by strong covalent bonds. It forms layers of hexagonal rings with no covalent bonds between layers. There are free electrons.

... they have strong covalent bonds and free electrons.